

Nitrogen (21-0-0) feedings by plant

Apply as side dressing at least 6 inches to the side of the plants.

Type	Amount	When to apply	Additional feedings
Artichoke	1-2 teaspoons per plant	4 weeks after emergence	every 4 weeks
Beans	NONE		
Beets	1/2 cup per 10 foot row	6 weeks after emergence	no
Broccoli	1 cup per 10 foot row	4 weeks after transplant	1/4 cup when head is size of quarter
Carrots	1 cup per 10 foot row	6 weeks after emergence	no
Cantaloupe	2 tablespoons per plant	After vines develop runners	no
Cauliflower	1 cup per 10 foot row	4 weeks after transplant	1/4 cup when head is size of quarter
Celery	1/2 cup per 10 foot row	4 weeks after transplant	no
Corn	1 cup per 10 foot row	When plants have 8-10 leaves	Add 1/2 cup per 10 foot row when silks appear
Cucumber	1/2 cup per 10 foot row	After vines develop runners	no
Eggplant	1 tablespoon per plant	4 weeks after transplant	8 weeks after transplant
Lettuce	1/2 cup per 10 foot row	4 weeks after transplant	no
Onions	1/2 cup per 10 foot row	4 weeks after transplant	Every 4 weeks until mid July.
Peas	NONE		
Peppers	1 tablespoon per plant	4 weeks after transplant	8 weeks after transplant
Potatoes	1/2 cup per 10 foot row	6 weeks after emergence	no
Pumpkin	2 tablespoons per plant	After vines develop runners	no
Radish	1/4 cup per 10 foot row	After emergence	no
Spinach	1/2 cup per 10 foot row	4 weeks after transplant	no
Squash	2 tablespoons per plant	After vines develop runners	no
Strawberry	1/2 cup per 10 foot row	At renovation	Again in late summer
Swiss Chard	1/2 cup per 10 foot row	4 weeks after transplant	no
Tomato	2 tablespoons per plant	4 weeks after transplant	8 weeks after transplant
Watermelon	2 tablespoons per plant	After vines develop runners	no